arXiv:1210.6397v1 [cond-mat.str-el] 23 Oct 2012

Nonadiabatic Time-Dependent Spin-Density Functional Theory for strongly correlated systems

Volodymyr Turkowski, Talat S. Rahman
Department of Physics and NanoScience and Technology Center,
University of Central Florida, Orlando, FL 32816
(Dated: October 25, 2012)

We propose a nonadiabatic time-dependent spin-density functional theory (TDSDFT) approach for studying the single-electron excited states and the ultrafast response of systems with strong electron correlations. The correlations are described by the correlation part of the nonadiabatic exchange-correlation (XC) kernel, which is constructed by using some exact results for the Hubbard model of strongly correlated electrons. We demonstrate that the corresponding nonadiabatic XC kernel reproduces main features of the spectrum of the Hubbard dimer and infinite-dimensional Hubbard model, some of which are impossible to obtain within the adiabatic approach. The theory may be applied for DFT study of strongly correlated electron systems in- and out-of-equilibrium, including the important case of nanostructures, for which it leads to a dramatic reduction of necessary computational power.

PACS numbers: 71.10.-w, 71.15.Mb, 71.27.+a

Introduction.–Description of the electronic properties of the systems that contain localized d- and f-orbitals remains one of the most important problems in condensed-matter physics for both extended and finite systems. The extended systems form a large class of materials with many exotic properties with in turn potentially many technological applications. Examples include cuprate high-temperature superconductors, heavy fermion materials and manganites. One can also expect many unusual properties in the much less explored cases of molecules and nanostructures, in which correlation effects may be even more enhanced owing to space constraint. Moreover, in modern nanotechnological applications, in which the distance between the atoms can be tuned (for example, by putting the atoms on a substrate), even s- and p-electron systems can find themselves in a strongly-correlated regime when the interatomic separation is large enough. Such systems show many surprises, like recently observed unusual antiferromagnetic ordering in small Fe chains,[1] metal-insulator transition in Au and Fe chains[2] etc. The excited states and the nonequilibrium properties of such systems, including the ultrafast (femto- and atto-second) response, are extremely important from the point of view of the modern trend of "smaller and faster." Correct understanding of the nanosystems and molecules with strong electron correlations may also shed light on the general properties of strongly correlated materials, including the local correlations and nonhomogeneous order in extended systems.

Most of the progress in studies of correlated systems has been made by using many-body approaches. Two of the most powerful of them are the Bethe ansatz in the 1D case and the dynamical mean-field theory (DMFT) approach, the latter of which is exact in the limit of infinite dimensions[3] and which appears to be also a good approximation in the 2D and 3D cases (reviews of both approaches are given in Refs. 4 and 5, correspondingly). These approaches have also been generalized to the nonequilibrium case,[6,7] allowing one to study the excitations and the nonlinear response of the corresponding systems. Since standard DFT approximations fail to describe properly most of the strongly-correlated effects, nowadays DMFT combined with DFT is used for this purpose.[8,9] This method allows one to describe the spectral, optical and magnetic properties of bulk and layered materials (see, for example, reviews 10,11). In DFT+DMFT, all properties of the "non-correlated" system (system geometry, bandstructure, etc.) are obtained within DFT (usually with the LDA or the GGA approximation), and the correlation effects are taken into account by solving the corresponding effective Hubbard model. Recently, it has been shown that a similar approach can be successfully applied to nanosystems.[12–18] Unfortunately, the combined DFT+DMFT calculations are computationally demanding even for the equilibrium case, especially for finite systems, which have a number of non-equivalent atoms.

For this reason it would be very useful to develop a TDDFT formalism with the corresponding XC potential that properly describes strongly correlated systems, including the excitations (in which case one needs to go beyond static DFT).[19] Some progress in this direction has already been made. For 1D systems, the corresponding adiabatic (static DFT) XC potentials were proposed in Refs. 20–23. In particular, in the last two papers a DFT approach based on the Bethe ansatz has been proposed. Recently, a combined adiabatic TDDFT-DMFT approach applicable in the 3D case has been proposed, in Ref. 24. In this approach, the XC potential is derived from the DMFT solution for the Hubbard model. The approaches were tested versus some exact solutions, and it was shown that they are successful in reproducing features of some of very important effects, including metal insulator transition and a temporal response of the systems. Unfortunately, some of the effects turned out not to be properly described. In particular, as was shown in

Ref. 24, the corresponding adiabatic potential fails to describe correctly the response of a finite Hubbard system in strongly correlated regime when the system is close to half-filling and when the local Coulomb repulsion U is large. One might expect that the main shortcomings come from the adiabatic approximation. Indeed, as one notes, the adiabatic XC kernel fails even to reproduce the correct number of the excited states of the system. For example, in the case of the Hubbard dimer, it is known that the local Coulomb repulsion – in addition to the bonding-anti-bonding orbital excitation – leads to a satellite level at finite U (see, e.g., Ref. 25). This extra state is related to the excited Hubbard "band" in the case of extended system (whence in fact comes the redistribution of the single-electron spectral weight to a higher energy (∼ U). It is also known from the DMFT solution that the Hubbard systems demonstrate an extra quasi-particle spectral weight at zero energy (chemical potential).[5] Although all these states are important in a strongly correlated regime, none can be reproduced with the adiabatic TDDFT. Indeed, in the adiabatic case the solution of the Casida equation[26] leads to a shift of the single-electron levels , it does not contain new states.

In this paper, we propose on the basis of some exact results for the Hubbard model a simple form of the nonadiabatic XC kernel which results in the single-electron spectrum of the Hubbard model that reproduces the main features of the spectra of both finite (dimer) and extended (infinite-dimensional) systems. This kernel can be easily implemented within the standard TDDFT codes for use in the strongly correlated case.

The Hubbard dimer.–In order to obtain the XC kernel $f_{XC\sigma\sigma'}(\mathbf{r},\mathbf{r}',\omega)$ for the Hubbard dimer, we map the eigenvalue equation derived from the dimer Green's function onto the corresponding TDDFT Casida eigenenergy equation.[26] The Casida equation for the dimer has the following general form:

$$\det\begin{bmatrix} \epsilon_0^2+2\epsilon_0 K_{\uparrow,\uparrow}(\omega)-\omega^2 & 2\epsilon_0 K_{\uparrow,\downarrow}(\omega) \\ 2\epsilon_0 K_{\downarrow,\uparrow}(\omega) & \epsilon_0^2+2\epsilon_0 K_{\downarrow,\downarrow}(\omega)-\omega^2 \end{bmatrix}=0, \tag{1}$$

where

$$K_{\sigma,\sigma'}(\omega)=\int\int \psi_g(\mathbf{r})\psi_u(\mathbf{r})\left(\frac{1}{|\mathbf{r}-\mathbf{r}'|}\right. \\ \left.+f_{XC\sigma,\sigma'}(\mathbf{r},\mathbf{r}',\omega)\right)\psi_g(\mathbf{r}')\psi_u(\mathbf{r}')d\mathbf{r}d\mathbf{r}' \tag{2}$$

(the first part being the Hartree term, and $f_{XC\sigma,\sigma'}(\mathbf{r},\mathbf{r}',\omega)$ Fourier transform of $\delta V_{XC\sigma}[n](\mathbf{r},t)/\delta n_{\sigma'}(\mathbf{r}',t')$ with respect to t-t'), $\epsilon_0=\epsilon_u-\epsilon_g$ is the excitation energy of the free electron, $\epsilon_{u,g}$ and $\psi_{u,g}(\mathbf{r})$ are the corresponding bonding- and anti-bonding energies and wave functions. This equation has two solutions

$$\omega^2=\epsilon_0(\epsilon_0+2(K_{\uparrow,\uparrow}\pm K_{\uparrow,\downarrow})), \tag{3}$$

where + corresponds to the singlet state, and − to the triplet one. Since the ground state of this system is singlet and the total spin of the isolated system is conserved, for definiteness we shall focus on the singlet state.

The dimer Green's function can be found from the exact solution for the Hubbard model with the Hamiltonian:

$$H=-t\sum_{i\neq j,\sigma}c^{\dagger}_{i\sigma}c_{j\sigma}+U\sum_i n_{i\uparrow}n_{i\downarrow}, \tag{4}$$

where $c^{\dagger}_{i\sigma}$ and $c_{j\sigma}$ are the creation and annihilation operator of electron with spin σ on site i, $n_{i\sigma}$ is the corresponding number of particle operator, and t is the hopping parameter. One can find the exact single particle dimer's Green's function, which has the following form in the singlet (or, more generally, the "non-magnetic") case:

$$\hat{G}^{-1}(\omega)=\begin{pmatrix} \omega-\Sigma_{11}(\omega) & t-\Sigma_{12}(\omega) \\ t-\Sigma_{21}(\omega) & \omega-\Sigma_{22}(\omega)\end{pmatrix}, \tag{5}$$

where the self-energies are

$$\Sigma_{11,\sigma\sigma'}=\Sigma_{22,\sigma\sigma'}=\delta_{\sigma\sigma'}\frac{U^2}{8}\left(\frac{1}{\omega-3t+i\delta}+\frac{1}{\omega+3t+i\delta}\right), \tag{6}$$

and

$$\Sigma_{12,\sigma\sigma'}=\Sigma_{21,\sigma\sigma'}=\delta_{\sigma\sigma'}\frac{U^2}{8}\left(\frac{1}{\omega-3t+i\delta}-\frac{1}{\omega+3t+i\delta}\right). \tag{7}$$

It is easy to show that, in this case, the corresponding eigenvalue problem has the following form:

$$t^2+\frac{U^2}{2}+\frac{U^2}{2}\frac{6t^2-U^2/4}{\omega^2-9t^2}=\omega^2. \tag{8}$$

Hence, one can obtain an equation for the XC kernel for the Hubbard dimer:

$$K_{\uparrow\uparrow}(\omega)+K_{\uparrow\downarrow}(\omega)=\frac{U^2}{4t}\left(1+\frac{6t^2-U^2/4}{\omega^2-9t^2}\right), \tag{9}$$

where $K_{\sigma\sigma'}(\omega)$ is defined in Eq. (2). Since we are interested in the contribution to the energy due to correlation effects, we assume that the Hartree ($\sim 1/|\mathbf{r}-\mathbf{r}'|$) and the exchange ($K_{\uparrow\uparrow}$) parts simply lead to the renormalization of the free-particle energy $\epsilon_0\to t$ in Eq. (1).

In this case, one can arrive at the following separable form of the correlation portion of the local XC the kernel, which follows from Eq. (9):

$$f_{C\uparrow,\downarrow}(\mathbf{r},\mathbf{r}',\omega)=\delta(\mathbf{r}-\mathbf{r}')\frac{\mathbf{U^2}}{\mathbf{4tA}}\left(\mathbf{1}+\frac{\mathbf{6t^2-U^2/4}}{\boldsymbol{\omega}^{\mathbf{2}}\mathbf{-9t^2}}\right), \tag{10}$$

were $A=\int\int|\psi_g|^2(\mathbf{r})|\psi_u|^2(\mathbf{r})d\mathbf{r}d\mathbf{r}'$. This kernel results in the exact excitation spectrum of the Hubbard dimer[25]:

$$\omega=\pm t\pm\sqrt{4t^2+\frac{U^2}{4}}. \tag{11}$$



FIG. 1: Exact and DMFT spectra for the Hubbard dimer as a function of U/t.

The spectra are presented in Fig.1a. As follows from this Figure, there are two extra states Hubbard satellite peaks with energy $\pm 3t$ at small U's. These states appear only at finite U, and come from the redistribution of single-electron spectral weight. It is easy to see that one cannot obtain these new states from Eq. (1) with the static XC kernel.

In the case of extended system, considered below, the DMFT approximation $\Sigma_{ij}(\omega) \simeq \delta_{ij}\Sigma(\omega)$ is valid. Even though in the case of the dimer this approximation is not sufficiently accurate, we present the corresponding expression for the kernel for the discussion below:

$$f_{C\uparrow,\downarrow}(\mathbf{r},\mathbf{r}',\omega) = \delta(\mathbf{r}-\mathbf{r}')\frac{U^2}{32tA}\left(8+\frac{72t^2-U^2}{\omega^2-9t^2} - \frac{9t^2U^2}{(\omega^2-9t^2)^2}\right). \quad (12)$$

As follows from the last two equations, they have the same expression at the high-frequency limit, but as $\omega \to 0$ they differ. The DMFT solution has an extra energy peak around zero energy, which is characteristic for this approach (Fig.1).

Extended systems.–In these cases, the DMFT with the local self-energy approximation can be applied. We shall use the following exact result for the high-frequency electron self-energy for the Hubbard model:

$$\Sigma_{ij}(\omega) = \delta_{ij}\left(U_i n_{i\downarrow} + \frac{U_i^2 n_{i\downarrow}(1-n_{i\downarrow})}{\omega}\right). \quad (13)$$

(see, for example Refs. 27,28). In the homogeneous case, one can construct the XC kernel by mapping the eigenvalue equation

$$\omega - \varepsilon_k - \Sigma(\omega) = 0, \quad (14)$$

or

$$\varepsilon_k^2 + 2\varepsilon_k Re[\Sigma(\omega)] + |\Sigma(\omega)|^2 = \omega^2, \quad (15)$$

onto the corresponding Casida equation. As in the dimer case one can find the equation which connects the XC kernel with the self-energy:

$$K_{\uparrow\downarrow}(\omega) \sim Re[\Sigma(\omega)]. \quad (16)$$

A more straightforward way to find the expression for the XC kernel is to compare the TDDFT and DMFT correlation energies:

$$E_C^{TDDFT} = \frac{1}{2}\int\int \delta n(\mathbf{r},t) f_{XC}(\mathbf{r},t;\mathbf{r}',t') \times \delta n(\mathbf{r}',t') d\mathbf{r}dtd\mathbf{r}'dt' \quad (17)$$

$$E_C^{DMFT} = \int\int \psi^*(\mathbf{r},t)\Sigma(t-t')\delta(\mathbf{r}-\mathbf{r}') \times \psi(\mathbf{r}',t') d\mathbf{r}dtd\mathbf{r}'dt' \quad (18)$$

Equations (13), (17) and (18), together with the result for the dimer, Eq. (10), can be used to construct the following "universal" (DMFT) function for the exchange part of the XC kernel for the extended systems:

$$f_{C\uparrow\downarrow}(\mathbf{r},\mathbf{r}',\omega) = \frac{\mathbf{U^2 F[n_0](r)}}{\mathbf{4t}}\delta(\mathbf{r}-\mathbf{r}')\frac{\mathbf{n_\downarrow(1-n_\downarrow)\omega}}{\mathbf{\omega^2 - B^2}}, \quad (19)$$

where $F[n_0](\mathbf{r})$ is a functional of the ground state density $n_0(\mathbf{r})$ (in general, of the spin parts) and $B \sim 3t$ in the case of dimer, while one can choose B^2 to be equal mean square of the kinetic energy ε_k^2, or more generally $B^2 = \alpha\varepsilon_k^2$, where $\alpha \sim 1$. $n_\downarrow$ is average per site number of the spin-down electrons. It is important to note that we do not include the contribution of the static Hartree term $Un_\downarrow$ to the self-energy, since it is canceled by the chemical potential at half-filling. With this XC kernel one can reproduce the main features of the spectrum of the infinite- dimensional Hubbard model: splitted (by energy U) bands and the zero energy quasi-particle peak, which disappears as U increases (Fig.2).

Conclusions.–As it is known from the DFT+DMFT study of real materials, the main effects of strong electron correlations can be described by local, temporally resolved electron interaction. Therefore, one can expect that a *spatially local, nonadiabatic* XC kernel is sufficient for description of the strongly correlated systems



FIG. 2: a) The DOS for the infinite-dimensional Hubabrd model with the hypercubic DOS $A(\epsilon) = (1/\sqrt{\pi}t^*)exp(-\epsilon^2/t^{*2})$ ($t^* = 2dt$ is the renormalized hopping, d - dimensionality of the system) and $F[n_0](\mathbf{r})$=1, A=1, $B = \varepsilon_{\mathbf{k}}$; b) the same in the low frequency case.

in the TDSDFT case. From our analysis above it follows that one can expect the following nonadiabatic XC kernel can be used to describe the main effects due to strong electron-electron correlations within TDDFT:

$$f_{XC\uparrow\downarrow}(\mathbf{r},\mathbf{r}',\omega) = \frac{\mathbf{U^2F[n_{\sigma0}](r)}}{\mathbf{4t}}\delta(\mathbf{r}-\mathbf{r}')\frac{\mathbf{n_{\downarrow}(1-n_{\downarrow})}\omega^{\alpha}}{\omega^{\mathbf{2}}-\mathbf{B^2}}, \qquad (20)$$

where B is a parameter proportional to the kinetic energy (B=3t in the case of dimer). The functional $F[n_{\sigma0}](\mathbf{r})$ is defined by the static (”non-correlated”, DFT) spin densities. In order to model strong on-site correlations it can be chosen (for spin-up electron) in the form $F[n_{\sigma0}](\mathbf{r}) = n_{\uparrow0}(\mathbf{r})/(n_{\uparrow0}(\mathbf{r}) + n_{\downarrow0}(\mathbf{r}))$ (equal 1/2 in the case of the dimer). Indeed , this function has the maximum when the corresponding site (space point with a high charge density) is occupied by spin-up electron only. The frequency power a in the numerator is expected to be between zero (in small systems, like dimer), and 1, as in the extended (DMFT) case. It is interesting to note that the case a=0 is in agreement with the exact asymptotic behavior of the XC kernel $f_{XC}(\omega \to \infty) \to A{+}B/\omega^2$, where A and B are constants.[29] The most essential features of the proposed XC kernel are spatial locality, proportionality to the local Coulomb repulsion, and the oscillating in time interaction of the electrons with opposite spin, $\sim \exp(\pm iBt)$, where the frequency of oscillations B is proportional to the hopping (kinetic) energy.

While further tests of the validity of the methodology presented here for larger scale systems, both in- and out-of-equilibrium is necessary, is on-going, the frequency dependence of the XC kernel in Eq. (20) is sufficient to describe the main features of single-orbital correlated systems. We expect the present approach to open the possibility of describing strongly correlated materials within the standard TDDFT framework, extending it beyond the metallic and semiconductor structures with plasmons, excitons and other excitations (see, e.g., Refs. 30,31).

Acknowledgements.–We thank Lyman Baker for critical reading of the manuscript and DOE for a partial support under grant DOE-DE-FG02-07ER46354.

[1] S. Loth, S. Baumann, C.P. Lutz, D. M. Eigler, A.J. Heinrich, Science **335**, 196 (2012).

[2] F. Fioravante, R. W. Nunes, Appl. Phys. Lett. **91**, 223115 (2007).

[3] W. Metzner and D. Vollhardt, Phys. Rev. Lett. **62**, 324 (1989).

[4] A.M. Tsvelick and P.W. Wiegmann, Adv. Phys. **32**, 453 (1983).

[5] A. Georges, G. Kotliar, W. Krauth, M.J. Rozenberg, Rev. Mod. Phys. **68**, 13 (1996).

[6] P. Mehta, N. Andrei, Phys. Rev. Lett. **96**, 216802 (2006).

[7] J.K. Freericks, V.M. Turkowski, V. Zlatic, Phys. Rev. Lett. **97**, 266408 (2006).

[8] V.I. Anisimov, A.I. Poteryaev, M.A. Korotin, A.O. Anokhin, G. Kotliar, J. Phys.: Condens. Matter **9**, 7359 (1997).

[9] A.I. Lichtenstein, M.I. Katsnelson, Phys. Rev. B **57**, 6884 (1998).

[10] G. Kotliar, S.Y. Savrasov, K. Haule V. S. Oudovenko, O. Parcollet,. C.A. Marianetti, Rev. Mod. Phys. **78**, 865 (2006).

[11] K. Held, I.A. Nekrasov, G. Keller, V. Eyert, N. Blumer, A.

McMahan, T. Pruschke. V.I. Anisimov, and D. Vollhardt, Phys. Stat. Sol. (b) **243**, 2599 (2006).

[12] S. Florens, Phys. Rev. Lett. **99**, 046402 (2007).

[13] D.W. Boukhvalov, L.I. Vergara, V.V. Dobrovitskii, M. I. Katsnelson, A. I. Lichtenstein, P. Koegerler, J. L. Musfeldt, and B. N. Harmon, Phys. Rev. B **77**, 180402 (2008).

[14] V. Turkowski, A. Kabir, N. Nayyar and T. S Rahman, J. Phys.: Condens. Matter **22**, 462202 (2010).

[15] V. Turkowski, A. Kabir, N. Nayyar and T. S Rahman, J. Chem. Phys. **136**, 114108 (2012).

[16] D. Jacob, K. Haule, and G. Kotliar, Phys. Rev. B **82**, 195115 (2010) .

[17] N. Lin, C. A. Marianetti, A.J. Millis, D.R. Reichman, Phys. Rev. Lett. **106**, 096402 (2011).

[18] D. Zgid, G. K.-L. Chan, Journ. of Chem. Phys. **134**, 094115 (2011).

[19] E. Runge and E.K.U. Gross, Phys. Rev. Lett. **52**, 997 (1984).

[20] N.A. Lima, M. F. Silva, L. N. Oliveira, K. Capelle, Phys. Rev. Lett. **90**, 146402 (2003); N.A. Lima, L.N. Oliveira, and K. Capelle, Europhys. Lett. **60**, 601 (2002).

[21] G. Xianlong, M. Polini, M.P. Tossi, V.L. Campo, Jr., K. Capelle, M. Rigol, Phys. Rev. B **73**, 165120 (2006).

[22] C. Verdozzi, Phys. Rev. Lett. **101**, 166401 (2008).

[23] W. Li, G. Xianlong, C. Kollath, and M. Polini, Phys. Rev. B **78**, 195109 (2008).

[24] D. Karlsson, A. Privitera, and C. Verdozzi, Phys. Rev. Lett. **106**, 116401 (2011).

[25] J.M. Tomczak , PhD Thesis, Ecole Polytechnique, France, 2007.

[26] M.E. Casida, in Recent Developments and Applications of Modern Density Functional Theory, Ed. J.M. Seminario, PP.391-439 (Elsevier, Amsterdam, 1996).

[27] J.K. Freericks, V. Turkowski, Phys. Rev. B **80**, 115119 (2009).

[28] V. Turkowski, J.K. Freericks, Phys. Rev. B **77**, 205102 (2008).

[29] R. van Leeuwen, Int. J. of Mod. Phys. B **15**, 1969 (2001).

[30] V. Turkowski, A. Leonardo, C.A. Ullrich, Phys. Rev. B **79**, 233201 (2009).

[31] N. Nayyar, V. Turkowski and T.S. Rahman, Phys. Rev. Lett. **109**, 157404 (2012).